

PERIOD~9

## **MATHEMATICS**

CHAPTER NUMBER :~ 2 CHAPTER NAME :~ POLYNOMIALS SUB TOPIC :~ RECAPITULATION OF POLYNOMIAL

CHANGING YOUR TOMORROW

Website: www.odmegroup.org Email: info@odmps.org

#### Toll Free: 1800 120 2316

Sishu Vihar, Infocity Road, Patia, Bhubaneswar-751024

## PREVIOUS KNOWLEDGE TEST

FACTORISE (i) 27y<sup>3</sup> + 125z<sup>3</sup>

(ii) 64m<sup>3</sup> – 343n<sup>3</sup>



#### LEARNING OUTCOME:~

1.Students will learn

Polynomials in one variable ,Term,Coefficient,General form and types of polynomials

- 2. Students will learn about Zeroes of the Polynomials
- 3.Students will learn about Division of Polynomials Remainder theorem,
- Factorization of Polynomials (Factor Theorem)
- Factorization of Polynomials using Factor Theorem and splitting the middle term, Algebraic identities
- Application of Algebraic identities.



## LEARNING OUTCOME:~

- Students will learn
- g) Division of Polynomials
- h)Remainder theorem
- i) Factorization of Polynomials (Factor Theorem)
- j)Factorization of Polynomials using Factor Theorem and splitting the middle term
- k)Algebraic identities
- 1) Application of Algebraic identities.



Question 8.

Factorise each of the following (i) 8a<sup>3</sup> +b<sup>3</sup> + 12a<sup>2</sup>b+6ab<sup>2</sup> (ii) 8a<sup>3</sup> ~b<sup>3</sup>~12a<sup>2</sup>b+6ab<sup>2</sup> (iii) 27~125a<sup>3</sup> ~135a+225a<sup>2</sup> (iv) 64a<sup>3</sup> ~27b<sup>3</sup> ~144a<sup>2</sup>b + 108ab<sup>2</sup>

Solution: (i)  $8a^3 + b^3 + 12a^2b + 6ab^2$  $= (2a)^3 + (b)^3 + 6ab(2a + b)$  $= (2a)^3 + (b)^3 + 3(2a)(b)(2a + b)$  $= (2 a + b)^3$ [Using  $a^3 + b^3 + 3 ab(a + b) = (a + b)^3$ ] = (2a + b)(2a + b)(2a + b)(ii)  $8a^3 - b^3 - 12o^2b + 6ab^2$  $= (2a)^3 - (b)^3 - 3(2a)(b)(2a - b)$  $= (2a - b)^3$ [Using  $a^3 + b^3 + 3 ab(a + b) = (a + b)^3$ ] = (2a - b) (2a - b) (2a - b)



(iii)  $27 - 125a^3 - 135a + 225a^2$ =  $(3)^3 - (5a)^3 - 3(3)(5a)(3 - 5a)$ =  $(3 - 5a)^3$ [Using  $a^3 + b^3 + 3 ab(a + b) = (a + b)^3$ ] = (3 - 5a)(3 - 5a)(3 - 5a)

(iv)  $64a^3 \sim 27b^3 \sim 144a^2b + 108ab^2$ =  $(4a)^3 - (3b)^3 - 3(4a)(3b)(4a - 3b)$ =  $(4a - 3b)^3$ [Using  $a^3 - b^3 - 3ab(a - b) = (a - b)^3$ ] = (4a - 3b)(4a - 3b)(4a - 3b)



#### Question 9. Verify

(i) 
$$x^3 + y^3 = (x + y) \cdot (x^2 - xy + y^2)$$
  
(ii)  $x^3 - y^3 = (x - y) (x^2 + xy + y^2)$ 

Solution:

(i) :: 
$$(x + y)^3 = x^3 + y^3 + 3xy(x + y)$$
  
 $\Rightarrow (x + y)^3 - 3(x + y)(xy) = x^3 + y^3$   
 $\Rightarrow (x + y)[(x + y)2 - 3xy] = x^3 + y^3$   
 $\Rightarrow (x + y)(x^2 + y^2 - xy) = x^3 + y^3$   
Hence, verified.

(ii) :: 
$$(x - y)^3 = x^3 - y^3 - 3xy(x - y)$$
  
 $\Rightarrow (x - y)^3 + 3xy(x - y) = x^3 - y^3$   
 $\Rightarrow (x - y)[(x - y)^2 + 3xy)] = x^3 - y^3$   
 $\Rightarrow (x - y)(x^2 + y^2 + xy) = x^3 - y^3$   
Hence, verified.



Question 10. Factorise each of the following

(i)  $27y^3 + 125z^3$ (ii)  $64m^3 - 343n^3$ 

Solution:

(i) We know that  $x^{3} + y^{3} = (x + y)(x^{2} - xy + y^{2})$ We have,  $27y^{3} + 125z^{3} = (3y)^{3} + (5z)^{3}$   $= (3y + 5z)[(3y)^{2} - (3y)(5z) + (5z)^{2}]$  $= (3y + 5z)(9y^{2} - 15yz + 25z^{2})$ 

(ii) We know that  $x^3 - y^3 = (x - y)(x^2 + xy + y^2)$ We have,  $64m^3 - 343n^3 = (4m)^3 - (7n)^3$   $= (4m - 7n)[(4m)^2 + (4m)(7n) + (7n)^2]$  $= (4m - 7n)(16m^2 + 28mn + 49n^2)$ 



Question 11.

Factorise  $27x^3 + y^3 + z^3 \sim 9xyz$ .

Solution:

We have,  $27x^3 + y^3 + z^3 - 9xyz = (3x)^3 + (y)^3 + (z)^3 - 3(3x)(y)(z)$ Using the identity,  $x^3 + y^3 + z^3 - 3xyz = (x + y + z)(x^2 + y^2 + z^2 - xy - yz - zx)$ We have,  $(3x)^3 + (y)^3 + (z)^3 - 3(3x)(y)(z)$   $= (3x + y + z)[(3x)^3 + y^3 + z^3 - (3x \times y) - (y \times 2) - (z \times 3x)]$  $= (3x + y + z)(9x^2 + y^2 + z^2 - 3xy - yz - 3zx)$ 



#### Question 12.

Verify that  $x^3 + y^3 + z^3 - 3xyz = 12 (x + y + z)[(x - y)^2 + (y - z)^2 + (z - x)^2]$ 

Solution:

## R.H.S $= 12(x + y + z)[(x - y)^{2} + (y - z)^{2} + (z - x)^{2}]$ $= 12 (x + y + 2)[(x^{2} + y^{2} - 2xy) + (y^{2} + z^{2} - 2yz) + (z^{2} + x^{2} - 2zx)]$ $= 12 (x + y + 2)(x^{2} + y^{2} + y^{2} + z^{2} + z^{2} + z^{2} - 2xy - 2yz - 2zx)$ $= 12 (x + y + z)[2(x^{2} + y^{2} + z^{2} - xy - yz - zx)]$ $= 2 \times 12 \times (x + v + z)(x^{2} + v^{2} + z^{2} - xv - vz - zx)$ $= (x + y + z)(x^{2} + y^{2} + z^{2} - xy - yz - zx)$ $= x^{3} + v^{3} + z^{3} - 3xvz = L.H.S.$ Hence, verified.



Question 13.

If x + y + z = 0, show that  $x^3 + y^3 + z^3 = 3$  xyz.

Solution:

Since, x + y + z = 0  $\Rightarrow x + y = -z (x + y)^3 = (-z)^3$   $\Rightarrow x^3 + y^3 + 3xy(x + y) = -z^3$   $\Rightarrow x^3 + y^3 + 3xy(-z) = -z^3 [\because x + y = -z]$   $\Rightarrow x^3 + y^3 - 3xyz = -z^3$   $\Rightarrow x^3 + y^3 + z^3 = 3xyz$ Hence, if x + y + z = 0, then  $x^3 + y^3 + z^3 = 3xyz$ 



Question 14.

Without actually calculating the cubes, find the value of each of the following (i)  $(-12)^3 + (7)^3 + (5)^3$ (ii)  $(28)^3 + (-15)^3 + (-13)^3$ 



Solution:

```
(i) We have, (-12)^3 + (7)^3 + (5)^3
Let x = -12, y = 7 and z = 5.
Then, x + y + z = -12 + 7 + 5 = 0
We know that if x + y + z = 0, then, x^3 + y^3 + z^3 = 3xyz
\therefore (-12)^3 + (7)^3 + (5)^3 = 3[(-12)(7)(5)]
= 3[-420] = -1260
```

```
(ii) We have, (28)^3 + (-15)^3 + (-13)^3
Let x = 28, y = -15 and z = -13.
Then, x + y + z = 28 - 15 - 13 = 0
We know that if x + y + z = 0, then x<sup>3</sup> + y<sup>3</sup> + z<sup>3</sup> = 3xyz
\therefore (28)^3 + (-15)^3 + (-13)^3 = 3(28)(-15)(-13)
= 3(5460) = 16380
```



Question 15.

Give possible expressions for the length and breadth of each of the following rectangles, in which their areas are given

- (i) Area  $25a^2 35a + 12$
- (ii) Area  $35y^2 + 13y 12$

Solution:

Area of a rectangle = (Length) x (Breadth) (i)  $25a^2 - 35a + 12 = 25a^2 - 20a - 15a + 12 = 5a(5a - 4) - 3(5a - 4) = (5a - 4)(5a - 3)$ Thus, the possible length and breadth are (5a - 3) and (5a - 4).

(ii) 
$$35y^2 + 13y - 12 = 35y^2 + 28y - 15y - 12$$
  
=  $7y(5y + 4) - 3(5y + 4) = (5y + 4)(7y - 3)$   
Thus, the possible length and breadth are  $(7y - 3)$  and  $(5y + 4)$ .



Question 16.

What are the possible expressions for the dimensions of the cuboids whose volumes are given below? (i) Volume  $3x^2 - 12x$ (ii) Volume  $12ky^2 + 8ky - 20k$ 

Solution:

Volume of a cuboid = (Length) x (Breadth) x (Height) (i) We have,  $3x^2 - 12x = 3(x^2 - 4x)$ = 3 x x x (x - 4)

: The possible dimensions of the cuboid are 3, x and (x - 4).



#### Solution:

Volume of a cuboid = (Length) x (Breadth) x (Height)

```
(ii) We have, 12ky^2 + 8ky - 20k

= 4[3ky^2 + 2ky - 5k] = 4[k(3y^2 + 2y - 5)]

= 4 \times k \times (3y^2 + 2y - 5)

= 4k[3y^2 - 3y + 5y - 5]

= 4k[3y(y - 1) + 5(y - 1)]

= 4k[(3y + 5) \times (y - 1)]

= 4kx (3y + 5) \times (y - 1)

Thus, the possible dimensions of the cuboid are 4k, (3y + 5) and (y - 1).
```



"As great a genius as Archimedes could not invent analytical geometry, for the algebraic knowledge necessary for such as achievement was not available in his time..."

~ Nathan A Court



### Evaluation:~

a)Without calculating cubes find(-12)<sup>3</sup>+(5)<sup>3</sup>+(7)<sup>3</sup>.
b) Find 'k' if x+3 is a factor of 3x<sup>2</sup>+kx+6.
c) If a+b+c=0, prove that a<sup>3</sup>+b<sup>3</sup>+c<sup>3</sup>=3abc.



## HOMEWORK:-REVISE THE CHAPTER -2



#### <u>AHA:~</u>

- 1. a-b=5, ab=12 Find  $a^2+b^2$ .
- 2. If  $x^3+ax^2-bx+10$  is divisible by  $x^3-3x+2$  find 'a' and 'b'.



# THANKING YOU ODM EDUCATIONAL GROUP

