

Exercise

Date _____
Page _____

Chapter - 2 (Physical Quantities And Measurement)

True or false

- 1) a) S.I. unit of temperature is fahrenheit. (F)
- b) Every measurement involves two things - a number and a unit (T).
- c) Mass is the measure of quantity of matter. (T)
- d) The S.I. unit of ~~quantity~~^{time is} hour. (F)
- e) The area can be expressed as the product of lengths of two sides. (T)

2) Fill in the blanks:

- a) The S.I. Unit length is metre.
of time is second, of mass is kilogram.
- b) °C is the unit of temperature.
- c) 1 metric tonne = 1000 kg.

- d) The zero mark in Celsius thermometer is the melting point of ice.
- e) The thermometer used to measure the human body temperature is called the clinical thermometer.
- f) The normal temperature of human body is 37.00000 °C or 98.6 °F.
- 3) a) Length of a housing plot - iv) Measuring tape
- b) Breadth of a book - vi) Metre ruler
- c) Mass of an apple - ii) Beam balance
- d) Period of time for study - i) clock
- e) Temperature of a body - iii) Thermometer
- f) Surface area - Graph paper of a leaf

Q) a) The symbol of degree Celsius
is:

- i) $^{\circ}\text{C}$.

b) 10 mm is equal to:

- ii) 1 cm.

c) The amount of surface occupied by an object

- iii) area.

d) A metric ruler is graduated in:

- iii) mm

e) A thermometer is graduated in:

- iii) $^{\circ}\text{C}$

B) Measurement is a comparison of an unknown quantity with a known fixed quantity of the same kind.

The value obtained on measuring a quantity is called its magnitude.

The magnitude of a quantity is expressed as numbers in its unit.


2) Two Characteristics of a unit are :-

1. It should be of convenient size.
2. It must be universally accepted; i.e. its value must remain same at all places and at all times.
- 3) In our daily life we measure the following four basic physical quantities.

1. Length
2. Mass
3. time
4. Temperature

4) S.I. units of


<u>Quantity</u>	<u>S.I. unit</u>	<u>Symbol of S.I. unit</u>
i) mass	kilogram	kg
ii) length	metre	m
iii) time	second	s
iv) temperature	Kelvin	K



5) One metre is defined as the distance travelled by light in $\frac{1}{299,792,458}$ of a second
 Multiple of metre = kilometre
 Submultiple of metre = centimetre


- 6) a) 12 inch = 1 ft
- b) 1 ft = 30.48 cm
- c) 20cm = 0.2 m
- d) 4.2m = 420 cm
- e) 0.2 km = 200 m
- f) 0.2 cm = 2 mm
- g) 1 yard = 0.91 m

7a) To measure the length of a Pencil - Using a metrule, Place metrule rule with its marking close to the Object. Let p be a pencil. The end P of the pencil coincides with the



- Zero mark on the ruler -
The end A of the pencil is.
Read by keeping the eye at the
position 'B' vertically above
the end A; So the length of
Pencil is 4.3 cm.

- b) The ends of the ruler get
damaged with use and its zero
mark may not be visible. To
measure the length of an object
with such a ruler, the object is
placed close to a specific
markings on the ruler and
positions of both ends of the
object are read on the ruler.
The difference of the two
readings gives the length of
the object. In fig. the reading
on ruler at the end X is 1.0 cm
and at the end Y is 4.3 cm.
So the length of the rod XY
is $4.3 - 1.0 = 3.3$ cm.



8. Name the device which you will use to measure the perimeter of your Play ground. Describe in steps how you will use it.

Ans We will ~~use~~ use a measuring tape to measure the perimeter of an playground. To measure the length of playground the tape is spread along the length of the curved area.

- 9) The diagram below shows a stick placed along a metre RULER. The length of the stick is measured keeping the eye at Positions A, B and C.

Ans a) ~~the~~ Length of Stick PQ from Position A = 3.4 cm

Position B = 3.2 cm

Position C = 3.0 cm

No they are not same.

b) Which is the correct position of the eye? Write the correct length of the stick.



ans. - 'B' is the correct position of the eye.
Correct length of the stick $PQ = 3.2\text{cm}$

10) The mass of a body is the quantity of matter contained in it. The S.I. unit of mass is kilogram. In short form, it is written as kg.

In C.G.S. System, the unit of mass is gram. (symbol g).

In F.P.S. system, the unit of mass is pound (symbol lb)

11) a) ~~2500 kg~~ $2500\text{kg} = 2.5\text{ metric tonne}$.

$$1000\text{kg} = \frac{1}{1000}\text{ metric tonne}$$

$$1\text{kg} = \frac{1}{1000}\text{ metric tonne}$$

$$\therefore 2500\text{kg} = \frac{1}{1000} \times 2500 \text{ metric tonne}$$

$$\therefore 2500\text{kg} = 2.5 \text{ metric tonne}$$

b) $150\text{kg} = 1.5 \text{ quintal}$

$$100\text{kg} = 1 \text{ quintal}$$

$$1\text{kg} = \frac{1}{100} \text{ quintal}$$

$$150\text{kg} = \frac{1}{100} \times 150 \text{ quintal}$$

$$\therefore 150\text{kg} = 1.5 \text{ quintal}$$

c) $10 \text{ lb} = 4,53,59 \text{ kg}$
 $1 \text{ lb} = 453.59 \text{ g}$
 $= 453.59 \times \frac{1}{1000} \text{ kg}$
 $= 0.45359 \text{ kg}$

$(\therefore 1 \text{ kg} = 1000 \text{ g})$

$\therefore 10 \text{ lb} = 0.45359 \times 10 \text{ kg}$
 $= 4.5359 \text{ kg}$
 $\therefore 10 \text{ lb} = 4.5359 \text{ kg}$

d) $2500 \text{ g} = 2.5 \text{ kg}$
 $1000 \text{ g} = 1 \text{ kg}$

$1 \text{ g} = \frac{1}{1000} \text{ kg}$

$\therefore 2500 \text{ g} = \frac{1}{1000} \times 2500 \text{ kg} = 2.5 \text{ kg}$
 $\therefore 2500 \text{ g} = 2.5 \text{ kg}$


e) $0.01 \text{ kg} = 10 \text{ g}$

$1 \text{ kg} = 1000 \text{ g}$

$\therefore 0.01 \text{ kg} = \frac{1}{1000} \times 0.01 \text{ g}$
 $= 1000 \times \frac{1}{100} \text{ g} = 10 \text{ g}$
 $\therefore 0.01 \text{ kg} = 10 \text{ g}$

b. ans

SUPPORT TO HOLD THE BALANCE



f) $5 \text{ mg} = 5 \times 10^{-6} \text{ kg}$

$$5 \text{ mg} = \frac{5}{1000} \text{ g or } 5 \times 10^{-3} \text{ g}$$

$$\frac{5}{1000} \text{ g or } 5 \times 10^{-3} \text{ g} = \frac{5}{1000 \times 1000} \text{ on } 5 \times 10^{-4} \text{ kg}$$

12. ~~one~~ Instrument commonly used to measure the mass of a body. Is the beam balance.

When we hold up the balance, we observe that when there is nothing on either pan, the beam is horizontal. The body whose mass is to be measured is placed on the left pan. The standard weight are put on the right pan. They are so adjusted that the standard weights gives the mass of the given body.

B.

The mass of 1 litre of water at 4°C is taken as 1 kilogram

$$1 \text{ quintal} = 100 \text{ kg}$$

$$1 \text{ metric ton} = 10 \text{ quintal} = 1000 \text{ kg}$$

14. The S.I. unit of time is second. In short form we write it as 's'.

One Second is the time interval between the two consecutive ticks that you hear from pendulum wall clock.

$$1 \text{ min} = 60 \text{ s}$$

$$1 \text{ h} = 60 \text{ min} = 3600 \text{ s}$$

$$1 \text{ day} = 24 \text{ h} = 86400 \text{ s}$$

$$1 \text{ year} = 365 \text{ days} = 3.15 \times 10^7 \text{ s}$$

15. Two devices used to measure the time interval of an event are (a) STOPCLOCK and (b) STOPWATCH

16. 1) $3\text{ minute} = 180\text{ second}$

$$1\text{ minute} = 60\text{ second}$$

$$3\text{ minutes } 15\text{ second} = 60 \times 3 + 15 \\ = 180 + 15$$

$$= 195\text{ seconds}$$

2) $1\text{ minute} = 60\text{ second}$

$$2\text{ minutes} = 2 \times 60 = 120\text{ second} \dots \dots (1)$$

$$1\text{ hour } 3600\text{ second}$$

$$5\text{ hours } 3600 \times 5 = 18000\text{ second} \dots \dots (2)$$

$$5\text{ hours } 2\text{ minutes and } 5\text{ second}$$

$$= 18000 + 120 + 5 = 18125$$

seconds

17. Temperature measures the degree of coldness and hotness of a body.

18. The S.I. unit of temperature is Kelvin (symbol K).

Common unit of temperature is degree centigrade (symbol °C)

19. Clinical thermometer - diagram - Refer book (pg - 28) (P.T.O)

Q - 19



fig. (CLINICAL THERMOMETER)

20. 1) melting Ice = 0°C

2) Boiling water = 100°C

21. Doctors use a special thermometer called the Clinical thermometer for measuring the

temperature of the patient's body. This thermometer has markings from 35°C to 42°C . It has a slight bend or kink in the stem just above the bulb. This kink is called the constriction. This constriction prevents the mercury from falling back all by itself. The temperature of a healthy person is 37°C . This temperature is marked by a red arrow.

22. To measure the temperature of a patient's body its bulb is kept either below the tongue or under the arm's pit of the Patient for about a minute. Then the thermometer is taken out and its reading is noted, when the temperature of patient's body is above 37°C , he is said to suffer with fever.
23. No a clinical thermometer can be used to measure the temperature of boiling water. The reasons are
1. It has a very small range.
 2. It can break on cooling and on excess heating.
- 24) The total surface occupied by an object is called its area or surface area.

25. The S.I. unit of area is Square metre which in short form is written as m².

26-i) Square yard : One square yard is the area of a square of each side 0.836 metre.

$$\begin{aligned} \text{1 square yard} &= 1 \text{ yard} \times 1 \text{ yard} \\ &= 0.9144 \text{ m} \times 0.9144 \text{ m} \\ &= 0.836 \text{ m}^2 \text{ (or } 0.84 \text{ m}^2 \text{ nearly)} \end{aligned}$$

ii) hectare : One hectare is the area of a square of each side 100 metre. Thus,

$$1 \text{ hectare} = 100 \text{ metre} = 10000 \text{ 'metre}^2 \text{ (or } 10^4 \text{ m}^2)$$

iii) km² : One square kilometre is the area of a square of each side 1 kilometre. Thus,

$$\begin{aligned} 1 \text{ km}^2 &= 1 \text{ km} \times 1 \text{ km} = 10000 \text{ m}^2 \\ &= 10^4 \text{ m}^2 \end{aligned}$$

Q7 iv) $\text{cm}^2 : 1\text{cm}^2 = \left(\frac{1}{100}\text{m}\right) \times \left(\frac{1}{100}\text{m}\right)$
 $= \frac{1}{10000} \text{m}^2$

v) $\text{mm}^2 : 1\text{mm}^2 = \frac{1}{10000}$
 $= 10^{-4} \text{ m}^2$

2nd. The area of a square can be calculated by using the following formula =

1. Area of square of side 1
 $= \text{side} \times \text{side}$
 $= 1 \times 1 = 1$

The area of a leaf is obtained by using a graph paper. A graph paper has small squares of each side 1 mm. The area of each big square is 1 cm^2 .

Procedure place the leaf on graph paper. Draw its outline on the paper and remove it.

Now count the number of complete squares. To this add the ~~number~~ number of incomplete squares which are half or more than half. Ignores which are less than half. Thus

Approximate Area = (No. of Complete Squares + no. of half or more than half of Incomplete Squares) X area of one Square.

- K